## PROPOSED FLETC TRAINING CENTER WETLANDS RARE PLANT SURVEY

LOCATION: Cheltenham, Prince Georges Co., MD

DRAINAGE: Piscataway Creek

GEOLOGY: Upland - Pleistocene Brandywine formation

Lower creeks eroded into Miocene Calvert formation

DESCRIPTION OF WETLANDS: See maps for approximate location and extent of wetlands.

<u>WKS-A:</u> Isolated depression surrounded by scrub pine. Almost all grasses — Panicum dicotomiflorum, Andropogon virginicus, Aristida longispica- with a few Juncus effusus and Eupatorium hyssopifolium.

WKS-C: Young maple - sweet gum woods with Cinna, stilt grass, Lycopus virginicus, Aster lateriflorus, and Carex spp.

WKS-D: Shaded by a thin canopy of red maple with scattered highbush blueberry. Mostly grass - Calamagnostis cinnoides and Chasmanthium laxum, with two mosses - Sphagnum sp. and Polystrichum type-and scattered Lycopus virginicus, Aster lateriflorus, and Osmunda cinnamomea.

WETLAND B: Stream corridor. Upper part has red maple, sweet gum, and greenbrier. Wider stretches have stilt grass. Middle part, near road crossing, is more open, with some maple, sweet gum, willow, arrowwood viburnum, and alder. Between shrubs is a heavy ground cover of goldenrods, deer tongue grass (Panicum clandestinum) and vines - honeysuckle and grape, with a few Carex spp. and Polygonumspp. in the stream channel. Lower part has heavy shade of mostly red maple with no shrubs and a carpet of stilt grass and wood reed grass (Cinna arundinacia), and a few Onoclea sensibalis, Lycopus virginicus. Boehmeria cylindrica, and Carex spp. Near the lower end is a small open patch dominated by Carex spp. and Polygonum spp.

HF-A & B: A wedge between road and fence. Nearly level young maple – sweet gum woods with greater diversity of both shrubs and herbs, including Cornus ammomum, Betula nigra, Ilex verticillata, Alnus serrulata, Scirpus cyperinus, Onoclea sensibalis, Aster lateriflorus Juncus spp., Carex spp., grass spp., several Scutellaria lateriflora and Lobelia cardinalis, and a population of Galium trifidum. This area has the only good quality forested wetland on the survey area. The tip of the wedge is open, with deer tongue grass and goldenrods.

HF-C: Erosion gully in maple woods with arrowwood viburnum at edge, stilt grass and Carex spp.

HF-D: A partly cleared strip along the fence with maple, spicebush, stilt grass, Cinna, and Glyceria striata.

HIF-E: Upper part is open ditch with sweet gum saplings on edge and some Carex lurida, Polygonum punctatum, Erichtites hicracifolia, between heavy patches of deer tongue grass and fescue. The lower part is maple woods with stilt grass and Cinna.

RB-A: Isolated bowl in middle of grassy field, with one willow in middle, and remnants of dead Phragmites. Herbaceous species include Scirpus cyperinus, Juncus effusus, Eupatorium coelestinum, Epilobum coloratum, and Aster spp.

RB-B: Another high-quality wetland with some forested habitat similar to HF\_A & B, and also with some opener, usually wet areas carpeted with Ludwigia palustris. Other species include Rosa carolina, Cephalanthus occidentalis, Callitriche sp., Echinocloa muricata, Panicum rigidulum, and Leersia oryzoides. Eastward the edges revert to maple – stilt grass woods, and there is an alder thicket near the east fence.

RB-7: Eroded stream channel through maple - sweet gum woods.

RARE PLANTS: Two species of concern have been reported from near the site:


Polygonum densiflorum – Dense-flowered knotweed – Endangered Galium trifidum – Small bedstraw – Undetermined


Polygonum densiflorum usually grows in shallow water. There is no good habitat for this species on the site, and no large Polygonums were seen.


Galium trifidum is found in moist woods and edges. It is a northern species, at the southern edge of it's range in Maryland. A well-established population is growing on the site in the wetlands associated with Piscataway Creek. Plants are scattered in alien-free and not heavily shrubby red maple woods, which is also the area of the highest-quality wetlands on the site.

SURVEYED: 15 October, 2001, approximately 4 hours

gamet Ebert







## In Bud ## estimated ##   1-5 sq. meter   % Immature   Feeble   % In Flower   1-10   5-10 sq. meter   % Ist year   \$\frac{7}{27}\$ Norm.  ## Immature Fruit   11-50   1-100 sq. meter   % Mature (established)   Exception   % Mature fruit   51-100   100 sq. meter   % Senescent   % Dormant   1001-10,000   actual   area (if known)   10K+	BE 03 Ebert
State: MD County(ies): Prince Georges Date: Source Code: Fold Quad Marpin #: Date: Source Code: Full extent of EO known and mapped?	
Full extent of EQ known and mapped?	
Full extent of EO known and mapped?	
Seed dispersing   101-1000   100K   100H   100H   100H   100K   100H	
BIOLOGY   Final   Code   Comment Name:   G   O   O   Tr   Tr   Tr   Tr   Tr   Tr	
Element Name: Galvo trification L.  Phenology Population Size Ramets Genets Genets I sq meter % Seedlings Very 1 sq meter % I sq meter % Insurance Freely 1 sq meter % Insurance from Insu	
Phenology   Population Size   Population Area   Age Structure   Vigor	
Seedings	Occ.#: _
# In Leaf	<u>r                                      </u>
# In Bud	feeble
% In Flower	ole
% Immature Fruit	nal
% Mature fruit	
% Seed dispersing	
Age structure unknown area (if known)    10K+	igorous
Topographic position    Topographic position   Crest   Court	
omments on above: Probably at least 20 plants most leng past flowering, with new vegetal vidence of reproduction? Keyes no Explain: A few late fruits you of reproduction: Sexual asexual both not sexual asexual asexual both not sexual asexual both not sexual asexual both not sexual asexual asex	
vidence of reproduction?	
ype of reproduction:	tive shocts
ny symbiotic or parasitic relationships?  yes	
Sipersal	
Spect   Slope   Light   Topographic position   Mosture	
N	
HABITAT  Spect Slope Light Topographic position Mosture  N NE 0-3 % Open Crest Inunda  E NW 3-8 % Partial Upper Slope Satura  S SE 8-15 % Filtered Mid-Slope Moist  W SW 15-35 % Shaded Lower Slope Dry-M  Flat 35% - Vertical Shaded Dry M  Cross section of topography (habitat) / Include scale, direction, element position	
Slope  N NE	
N         NE         ✓ 0-3 %         Open         Crest         Inundation           E         NW         3-8 %         Partial         Upper Slope         ✓ Saturation           S         SE         8-15 %         ✓ Filtered         Mid-Slope         Moist           W         SW         15-35 %         Shaded         Lower Slope         Dry-M           X         Flat         35% - Vertical         ✓ Bottom         Dry (X           levation:         2 1 5 0 / ft to         ft         Cross section of topography (habitat) / Include scale, direction, element position	
E NW 3-8 % Partial Upper Slope Satura  S SE 8-15 % Filtered Mid-Slope Moist  W SW 15-35 % Shaded Lower Slope Dry-M  Flat 35% - Vertical Shaded Dry (X  levation: 5 15 0 / ft to ft  Cross section of topography (habitat) / Include scale, direction, element position	dated (Hydric)
S         SE         8-15 %         ★ Filtered         Mid-Slope         Moist           W         SW         15-35 %         Shaded         Lower Slope         Dry-M           ★ Flat         35% - Vertical         ★ Bottom         Dry (X           evation:         - 15 € € / It to	
wsw15-35 %ShadedLower SlopeDry-M  X Flat35% - VerticalX BottomDry (X  evation: = 150	
X Flat 35% - Vertical X Bottom Dry (X evation: 5150 ft to ft  Cross section of topography (habitat) / Include scale, direction, element position	
levation: = 150 / ft toft  Cross section of topography (habitat) / Include scale, direction, element position	
Cross section of topography (habitat) / Include scale, direction, element position	xenc)
Cross section of topography (habitat) / Include scale, direction, element position	
Gitrifidum scattered in more open areas  Gitrifidum scattered in more open areas in interior of wetland, on slightly raised temps	
in interior of wetland, on elightly raised temps	
	,
Company france	
Dier Open France  Gross Solidage School	

Not land at book 2 dgs or floody kin

nabilial (Continued)

Associated natural community/plant community: Wet young maple-sweet gum woods - fairly open under
· · · · · · · · · · · · · · · · · · ·
Associated plant species Acer robrum Liquidambar styraciflua, Alnus serrulate
Associated plant species Acer robrum Liquidam Bar styreetings Pancum dicatement
quercus palustris Cinna arundinacia scirpus experimes, somtellaria lateriflora
Associated plant species Acer roution English Scirpus experious, Panicom dicetements Suercus palostris Cinna arundinacia Scirpus experious, Panicom dicetement Aster lateriflorus, Onoclea sensibalis, Juncus spp scutellaria lateriflora Carex spp.
Soil name(s)/Substrate:
Estimated # of hectares of potential habitat in the immediate area:
IDENTIFICATION
Potograph taken?yes _x_no
Specimen taken? X yes no If yes, give collector, collection # and repository: 50 reyor has very poor specimen
Do other members of this genus co-occur at this site? X yes no If yes, complet below:
List: Possible G. tincterium  (could only find a few date-season inflorescene's  Hybridization? yes no mostly second effect translated y uniform, longer, no recover leaves.  Identification problems? yes no Explain: ID based on consistantly y uniform, longer, no recover leaves.
Hybridization? yes no
Identification problems? X yes no Explain: ID based on consistantly 4 uniform, longer, noticed to the same for the same to
CONSERVATION Pedicels James July 1
Owner protecting EO? yes no Unknown
Owner and cords.
Little Control of the
Threats to EO: Change in hydrology  Conservation/management needs: Keep Floedplain wetland in fact.
Conservation/management needs: Keep Floedplain wer takes mendy preserved ??
Areas upstream and down strength attempt pro
Research needs:
Data security?yesno Explain:
SUMMARY
EO Quality: (i.e., How representative is this occurence? Consider the size and productivity of the population and the vitality and vigor of the individuals.)
D. Boor
A. Excellent B. Good C. Marginal D. Poor
Comments:(
EO Condition: (i.e., is the habitat supporting the EO pristine or degraded? Is there a potential for the habitat to recover from disturbances?
A - Excellent B - Good C - Marginal D - Poor
Comments:
EO Viability: (i.e., what are the long-term prospects for continued existence of this occurrence at the indicated level of quality?)
A - Excellent B - Good C - Marginal D - Poor
comments: bood if whole area is preserved
EO Defensibility: (i.e., can this occurrence be protected from intrinsic human factors?)
A - Excellent B - Good C - Margina D - Poor
Comments:
EO Rank: (i.e., a summary of all factors listed above) A B C D
EO Rank: (i.e., a summary of all factors holds assets)
Comments: Nice habitat volnirable